LEGACY HEALTH	Legacy Day Treatment Unit Provider's Orders Adult Ambulatory Infusion Order DENOSUMAB (PROLIA) OSTEOPOROSIS	Patient Name: Date of Birth: Med. Rec. No (TVC MRN Only):			
ALL ORDERS MUST BE MARKED IN INK WITH A CHECKMARK (✓) TO BE ACTIVE					
Anticipated Start Date: Patient to follow up with provider on date: ***This plan will expire after 365 days, unless otherwise specified below***					
	kpire:kg Height:	cm			

GUIDELINES FOR PRESCRIBING:

Diagnosis: _____

- 1. Send FACE SHEET, INSURANCE CARD and most recent provider chart or progress note.
- 2. Risk versus benefit regarding osteonecrosis of the jaw and hip fracture must be discussed prior to treatment.

Diagnosis Code: _____


- 3. Hypocalcemia must be corrected before initiation of therapy. The corrected calcium level should be greater than or equal to 8.4 mg/dL.
- 4. All patients should be prescribed daily calcium and vitamin D supplementation.
 - a. Recommended dosing: calcium 1000 mg and vitamin D 400 IU daily
- 5. A complete metabolic panel must be drawn within 60 days prior to starting treatment.
- 6. In patients with severe renal impairment (creatinine clearance less than 30 mL/min), high risk of hypocalcemia, disturbances of mineral metabolism (e.g. hypoparathyroidism, thyroid surgery, parathyroid surgery, malabsorption syndromes, excision of small intestines) recommend clinical monitoring of calcium, magnesium and phosphorus levels within 14 days of Prolia injection.
- 7. Pregnancy must be ruled out prior to administration. Perform pregnancy testing in all females of reproductive potential prior to administration of Prolia. It is the responsibility of the ordering provider to determine necessity and obtain results, if indicated. This is not a hold parameter for infusion staff.
- 8. Provider confirms that the patient has had a recent oral or dental evaluation and/or has no contraindications to therapy related to dental issues prior to initiating therapy.

LABS TO BE DRAWN (orders must be placed in TVC Epic by ordering provider if TVC provider):

CMP, Routine, every 6 months prior to Prolia dose

MEDICATIONS:

 denosumab (PROLIA) 60 mg (1 mL) SUBCUTANEOUSLY, every 6 months (26 weeks) for 2 treatments Administer injection into upper arm, upper thigh, or abdomen



ALL ORDERS MUST BE MARKED IN INK WITH A CHECKMARK (✓) TO BE ACTIVE

NURSING ORDERS (TREATMENT PARAMETERS):

- 1. Review previous creatinine clearance, serum calcium and albumin. If no results in past 60 days order CMP.
- 2. Treatment parameters, ONCE: Hold and notify MD for corrected calcium LESS than 8.4 mg/dL.
- 3. If corrected calcium is between 8.4 and 8.8 or creatinine clearance LESS than 30 mL/min review home medication for calcium and vitamin D supplementation. If patient is not on these agents, notify provider.
- 4. Do not hold treatment for CrCl <30 mL/min
- 5. Assess for new or unusual thigh, hip, groin, or jaw pain. Inform provider if positive findings or if patient is anticipating invasive dental work.
- 6. Remind patient to take calcium and vitamin D supplements, as directed by prescribing provider.

Please check the appropriate box for the patient's preferred clinic location:

Legacy Day Treatment Unit –

The Vancouver Clinic Building A department of Salmon Creek Medical Center 700 NE 87th Avenue, Suite 360 Vancouver, WA 98664 Phone number: 360-896-7070 Fax number: 360-487-5773

□ Legacy Salmon Creek Day Treatment Unit

Legacy Salmon Creek Medical Center 2121 NE 139th Street, Suite 110 Vancouver, WA 98686 Phone number: 360-487-1750 Fax number: 360-487-5773

□ Legacy Emanuel Day Treatment Unit

A department of Emanuel Medical Center 501 N Graham Street, Suite 540 Portland, OR 97227 Phone number: 503-413-4608 Fax number: 503-413-4887

□ Legacy STEPS Clinic

A department of Silverton Medical Center Legacy Woodburn Health Center 1475 Mt Hood Ave Woodburn, OR 97071 Phone number: 503-982-1280 Fax number: 503-225-8723

Provider signature:	Date/Time:		
Printed Name:	Phone:	Fax:	
Organization/Department:			