



Legacy Day Treatment Unit Provider's Orders

Adult Ambulatory Infusion Order
INFLIXIMAB-dyyb (INFLECTRA)
& INFLIXIMAB (REMICADE)

Patient Name:

Date of Birth:

Med. Rec. No (TVC MRN Only):

ALL ORDERS MUST BE MARKED IN INK WITH A CHECKMARK (✓) TO BE ACTIVE

Anticipated Start Date: _____ Patient to follow up with provider on date: _____

This plan will expire after 365 days, unless otherwise specified below

Orders expire: _____

Weight: _____ kg Height: _____ cm

Allergies: _____

Diagnosis: _____ Diagnosis Code: _____

GUIDELINES FOR PRESCRIBING:

1. Send FACE SHEET, INSURANCE CARD and most recent provider chart or progress note.
2. A Tuberculin test (PPD) or QuantiFERON Gold must have been read as negative prior to initiation
If the QuantiFERON Gold is indeterminate a CHEST X-ray should be performed to rule out infection
3. Hepatitis B (Hep B surface antigen AND core antibody) screening must be completed prior to initiation of therapy and the patient should not be infected
4. Patient should not have an active ongoing infection, signs or symptoms of malignancy, or moderate to severe heart failure at the onset of inFLIXimab therapy. Baseline liver function tests should be normal
5. Patients should have regular monitoring for TB, Hepatitis B, infection, malignancy, and liver abnormalities throughout therapy

PRE-SCREENING: (Results must be available prior to initiation of therapy)

• Hepatitis B Surface AG	Result Date: _____	<input type="checkbox"/> Positive	<input type="checkbox"/> Negative
• Hepatitis B Core AB Qual	Result Date: _____	<input type="checkbox"/> Positive	<input type="checkbox"/> Negative
• Tuberculin Test	Result Date: _____	<input type="checkbox"/> Positive	<input type="checkbox"/> Negative
• QuantiFERON Gold Test	Result Date: _____	<input type="checkbox"/> Positive	<input type="checkbox"/> Negative
• Chest X-ray (if QuantiFERON Gold indeterminate)	Result Date: _____	<input type="checkbox"/> Positive	<input type="checkbox"/> Negative
• Baseline CMP/LFT	Result Date: _____	<input type="checkbox"/> WNL	<input type="checkbox"/> Negative

LABS TO BE DRAWN (orders must be placed in TVC Epic by ordering provider if TVC provider):

Complete Metabolic Panel, Routine, ONCE, every _____(visit)(days)(weeks)(months)- **Circle one**

CBC with differential, Routine, ONCE, every _____(visit)(days)(weeks)(months)- **Circle one**

Other: _____



Legacy Day Treatment Unit Provider's Orders

Adult Ambulatory Infusion Order
INFLIXIMAB-dyyb (INFLECTRA)
& INFLIXIMAB (REMICADE)

Patient Name:

Date of Birth:

Med. Rec. No (TVC MRN Only):

ALL ORDERS MUST BE MARKED IN INK WITH A CHECKMARK (✓) TO BE ACTIVE

PRE-MEDICATIONS:

- acetaminophen (TYLENOL) tablet, 650 mg, oral, ONCE, every visit
- diphenhydramine (BENADRYL) tablet, 50 mg, oral, ONCE, every visit
(Give either cetirizine or diphenhydramine, not both.)
- cetirizine (ZYRTEC) tablet, 10 mg, oral, ONCE, every visit
(Give either cetirizine or diphenhydramine, not both.)
- methylPREDNISolone sodium succinate (SOLU-MEDROL), 40 mg, IV, ONCE, every visit
(Choose if patient has required IV steroids for a reaction during prior TNF-alpha inhibitor infusion.)

MEDICATIONS: (must check one):

Biosimilar selection (must check one) – applies to all orders below

- INFLECTRA (inFLIXimab-dyyb) ****formulary agent****
- REMICADE (inFLIXimab) *Restricted ONLY to existing REMICADE patients for continuing therapy, or patients whose insurance will only cover REMICADE*
- RENFLEXIS (inFLIXimab-abda)
- AVSOLA (inFLIXimab-axxq)

Dose: (Pharmacist will use most recent weight and round dose to the nearest 100 mg vial – dose selected at initiation will be continued, unless new orders are received specifying a dose modification)

- 3 mg/kg in NaCl 0.9% IV, ONCE, every visit
- 5 mg/kg in NaCl 0.9% IV, ONCE, every visit
- 10 mg/kg in NaCl 0.9% IV, ONCE, every visit
- _____ mg/kg in NaCl 0.9% IV, ONCE, every visit
- _____ mg in NaCl 0.9% IV, ONCE, every visit

Interval:

- Once
- Three doses at Week 0, 2, and 6, then every _____ weeks
- Every _____ weeks
- Other: _____



Legacy Day Treatment Unit Provider's Orders

Adult Ambulatory Infusion Order
INFLIXIMAB-dyyb (INFLECTRA)
& INFLIXIMAB (REMICADE)

Patient Name:

Date of Birth:

Med. Rec. No (TVC MRN Only):

ALL ORDERS MUST BE MARKED IN INK WITH A CHECKMARK (✓) TO BE ACTIVE

AS NEEDED MEDICATIONS:

- acetaminophen 650 mg oral, EVERY 4 HOURS AS NEEDED for hypersensitivity or infusion reaction, chills, or malaise
- diphenhydrAMINE 25 mg oral, may repeat x 1 EVERY 4 HOURS AS NEEDED for itching
- NaCl 0.9% 500 mL IV, AS NEEDED, ONCE, infusion tolerability. Give concurrently with inFLIXimab/inFLIXimab biosimilars

NURSING ORDERS (TREATMENT PARAMETERS):

- Hold treatment and contact provider if Hepatitis B surface antigen or core antibody total test result is positive, TB test result is positive, or if screening has not been performed.
- Hold infusion and contact provider if patient has signs or symptoms of infection.
- Monitor and record vital signs, tolerance and presence of infusion-related reactions prior to infusion and at the end of the infusion.
- Monitor vital signs every 15 minutes for the first hour, then every 30 minutes for remainder of infusion. Stop the infusion immediately if a reaction occurs.
- For 1st infusion or for any patient with history of infusion-related adverse reactions, monitor patient for 30 minutes after completion of treatment.
- Manage line per LH policy 904.4007 IV Catheter Insertion (Peripheral) and LH 904.4004 IV Access: Central Catheters
- Manage hypersensitivity reactions per LH 906.6606

HYPERSensitivity MEDICATIONS: Refer to LH policy 906.6606 Initiation of Emergency Measures for Adult Oncology, Radiation Oncology and Infusion Clinic Patients

- hydrocortisone 100 mg IV, AS NEEDED x1 dose for hypersensitivity reaction
- diphenhydrAMINE 25-50 mg IV, EVERY 2 HOURS AS NEEDED for hypersensitivity reaction (Max dose: 50 mg)
- famotidine 20 mg IV, AS NEEDED x1 dose for hypersensitivity reaction
- EPINEPPhrine 0.3 mg IM, AS NEEDED x1 dose for hypersensitivity reaction
- naloxone (Narcan) 0.4 mg IV, AS NEEDED x1 dose for diminished respiratory rate
- Nursing communication order, every visit: Please follow treatment algorithm for acute infusion reaction



Legacy Day Treatment Unit Provider's Orders

Adult Ambulatory Infusion Order
INFLIXIMAB-dyyb (INFLECTRA)
& INFLIXIMAB (REMICADE)

Patient Name:

Date of Birth:

Med. Rec. No (TVC MRN Only):

ALL ORDERS MUST BE MARKED IN INK WITH A CHECKMARK (✓) TO BE ACTIVE

Please check the appropriate box for the patient's preferred clinic location:

**Legacy Day Treatment Unit –
The Vancouver Clinic Building**
A department of Salmon Creek Medical Center
700 NE 87th Avenue, Suite 360
Vancouver, WA 98664
Phone number: 360-896-7070
Fax number: 360-487-5773

Legacy Emanuel Day Treatment Unit
A department of Emanuel Medical Center
501 N Graham Street, Suite 540
Portland, OR 97227
Phone number: 503-413-4608
Fax number: 503-413-4887

Legacy Salmon Creek Day Treatment Unit
Legacy Salmon Creek Medical Center
2121 NE 139th Street, Suite 110
Vancouver, WA 98686
Phone number: 360-487-1750
Fax number: 360-487-5773

Legacy STEPS Clinic
A department of Silverton Medical Center
Legacy Woodburn Health Center
1475 Mt Hood Ave
Woodburn, OR 97071
Phone number: 503-982-1280
Fax number: 503-225-8723

Provider signature: _____ **Date/Time:** _____

Printed Name: _____ **Phone:** _____ **Fax:** _____

Organization/Department: _____