

Legacy Day Treatment Unit Provider's Orders

Adult Ambulatory Infusion Order RITUXIMAB (RITUXAN) FOR NON-ONCOLOGY INDICATIONS

Patient Name:
Date of Birth:
Med. Rec. No (TVC MRN Only):

ALL ORDERS MUST BE MARKED IN INK WITH A CHECKMARK (✓) TO BE ACTIVE

This p		ter 365 days, unle	tient to follow up with provider on date: ess otherwise specified below	
Weight:	kg	Height:	cm	
Allergie	s:			
Diagnos	is:		Diagnosis Code:	
GUIDEL	INES FOR PRESO	CRIBING:		
2. H	lepatitis B (Hep B		ARD and most recent provider chart or progress note. nd core antibody) screening must be completed prior to initiate infected	ion of
PRE-SC	REENING: (Resu	lts must be availa	able prior to initiation of therapy)	
	-		Date: Desitive / Negative Ult Date: Positive / Negative	
LABS TO	O BE DRAWN (or	ders must be plac	ced in TVC Epic by ordering provider if TVC provider):	
	BC with differentia	al, Routine, every	ery(visit)(days)(weeks)(months) - Circle one(visit)(days)(weeks)(months) - Circle one	
	cetaminophen (TY 650 mg 325 mg	minister 30 minutes 'LENOL) tablet, ora	•	
	☐ 25 mg ☐ 50 mg	(BENADRYL) table		dod)
	☐ 10 mg		y visit (Choose as alternative to diphenhydramine, if need ate (SOLU-MEDROL) IV, every visit	iea)



Legacy Day Treatment Unit Provider's Orders

Adult Ambulatory Infusion Order RITUXIMAB (RITUXAN) FOR NON-ONCOLOGY INDICATIONS

Patient Name:	
Date of Birth:	
Med. Rec. No (TVC MRN Only):	

ALL ORDERS MUST BE MARKED IN INK WITH A CHECKMARK (✓) TO BE ACTIVE

	CATIONS: (must check one): se: (Pharmacist will use most recent weight at initiation and round dose to the nearest whole vial)
	riTUXimab (RITUXAN) 375 mg/m2 = mg IV in NaCl 0.9% to a final concentration of 2 mg/mL. First infusion or prior infusion reactions: start at 50 mg/hr x 30 min, then may increase by 50 mg/hr every 30 minutes if tolerated (NTE 400 mg/hr). Subsequent infusions if no reactions infuse rituximab at 100 mg/hr for the first 30 min. If no infusion related reactions are seen, increase rate by 100 mg/hour every 30 minutes to a maximum of 400 mg/hr
	riTUXimab (RITUXAN) mg IV in NaCl 0.9% to a final concentration 2 mg/mL. First infusion or prior infusion reactions: start at 50 mg/hr x 30 min, then may increase by 50 mg/hr every 30 minutes if tolerated (NTE 400 mg/hr). Subsequent infusions if no reactions infuse rituximab at 100 mg/hr for the first 30 min. If no infusion related reactions are seen, increase rate by 100 mg/hour every 30 minutes to a maximum of 400 mg/hr
Interv	al:
	Once Every 2 weeks x 2 doses Every weeks x doses Weekly x 4 doses Other

AS NEEDED MEDICATIONS:

- acetaminophen 650 mg oral, EVERY 4 HOURS AS NEEDED for fever, headache or pain
- diphenhydrAMINE 25 mg oral, EVERY 4 HOURS AS NEEDED for itching
- meperidine 25-50 mg IV, EVERY 2 HOURS AS NEEDED (NTE 50 mg/hr) for rigors in the absence of hypotension

NURSING ORDERS (TREATMENT PARAMETERS):

- 1. Vital signs, every visit: Monitor and record vital signs, tolerance, and presence of infusion-related reactions prior to infusion and at the end of infusion.
- 2. Nursing communication order, every visit: Manage line per LH policy 904.4007 IV Catheter Insertion (Peripheral) and LH 904.4004 IV Access: Central Catheters.
- 3. Nursing communication orders, every visit: Manage hypersensitivity reactions per LH 906.6606.



Legacy Day Treatment Unit Provider's Orders

Adult Ambulatory Infusion Order RITUXIMAB (RITUXAN) FOR NON-ONCOLOGY INDICATIONS

Patient Name:		
Date of Birth:		

Med. Rec. No (TVC MRN Only):

ALL ORDERS MUST BE MARKED IN INK WITH A CHECKMARK (✓) TO BE ACTIVE

HYPERSENSITIVITY MEDICATIONS: Refer to LH policy 906.6606 Initiation of Emergency Measures for Adult Oncology, Radiation Oncology and Infusion Clinic Patients

- 1. hydrocortisone 100 mg IV, AS NEEDED x1 dose for hypersensitivity reaction
- 2. diphenhydrAMINE 25-50 mg IV, EVERY 2 HOURS AS NEEDED for hypersensitivity reaction (Max dose: 50 mg)
- 3. famotidine 20 mg IV, AS NEEDED x1 dose for hypersensitivity reaction
- 4. EPINEPHrine 0.3 mg IM, AS NEEDED x1 dose for hypersensitivity reaction
- 5. naloxone (Narcan) 0.4 mg IV, AS NEEDED x1 dose for diminished respiratory rate
- 6. sodium chloride 0.9% 1000 mL IV, AS NEEDED x 1 dose for alteration in hemodynamic status
- 7. Nursing communication order, every visit: Please follow treatment algorithm for acute infusion reaction.

Please check the appropriate box for the patient's preferred clinic location: ☐ Legacy Day Treatment Unit – ☐ Legacy Emanuel Day Treatment Unit The Vancouver Clinic Building A department of Emanuel Medical Center A department of Salmon Creek Medical Center 501 N Graham Street, Suite 540 700 NE 87th Avenue, Suite 360 Portland, OR 97227 Vancouver, WA 98664 Phone number: 503-413-4608 Fax number: 503-413-4887 Phone number: 360-896-7070 Fax number: 360-487-5773 ☐ Legacy Silverton STEPS Clinic ☐ Legacy Salmon Creek Day Treatment Unit Legacy Salmon Creek Medical Center Legacy Silverton Medical Center 2121 NE 139th Street, Suite 110 342 Fairview Street Vancouver, WA 98686 Silverton, OR 97381 Phone number: 360-487-1750 Phone number: 503-873-1670 Fax number: 360-487-5773 Fax number: 503-874-2483 ☐ Legacy Woodburn STEPS Clinic A department of Silverton Medical Center Legacy Woodburn Health Center 1475 Mt Hood Ave Woodburn, OR 97071 Phone number: 503-982-1280 Fax number: 503-225-8723 Date/Time: ____ Provider signature: Phone: _____ Fax: ____ Printed Name: _____ Organization/Department: _____